

C-TOB-D

5ml Approx.
Eye Drops

Tobramycin 3 mg
Dexamethasone 1 mg
Sterile Ophthalmic Suspension
(USP Specification)

COMPOSITION:

Each ml contains:
Tobramycin 3mg
Dexamethasone 1mg
(USP Specification)

INDICATIONS:

C-TOB-D ophthalmic suspension is indicated for the reduction of ocular inflammation and prophylaxis of infection due to susceptible organisms.

CONTRA-INDICATIONS:

Epithelial herpes simplex keratitis (dendritic keratitis), vaccinia, varicella and many other viral disease of the cornea and conjunctiva. Mycobacterial infection of the eye. Fungal diseases of ocular structures. Hypersensitivity to any component of the medication. The use of this combination is always contraindicated after uncomplicated removal of a corneal foreign body. Topical corticosteroids should not be used in the treatment of mechanical lacerations and abrasions of the eye. They delay healing and promotes the development and spread of infection.

WARNINGS:

Prolonged use of topical ophthalmic steroids may result in glaucoma, with damage to the optic nerve, defects in visual acuity and fields of vision and posterior subcapsular cataract formation. IF THESE PRODUCTS ARE USED FOR 10 DAYS OR LONGER INTRA OCULAR PRESSURE SHOULD BE ROUTINELY MONITORED EVEN THOUGHT IT MAY BE DIFFICULT IN CHILDREN AND UN CO-OPERATIVE PATIENT. In those diseases causing thinning of the cornea or sclera, perforations have been known to occur with the use of topical steroids. Corneal ulceration may be aggravated when corticosteroids are applied. It is important that corneal ulcers are correctly designed before treatment with corticosteroids is initiated. In acute purulent conditions of the eye, steroids may mask progression of infection until sight is lost or enhance existing infection. Not for injection into the eye. This product contains benzalkonium chloride and is not recommended for use whilst soft contact lenses are being worn. Sensitivity to topically applied amino glycosides may occur in some patients. If a sensitivity reaction does occur, discontinue use.

Dosage and administrations For Use:

C-TOB-D ophthalmic suspension: Instill 1-2 drops into the effective eye/eyes 3 to 4 times a day or as directed by the physician.

SIDE-EFFECTS:

The most frequent side effects to topical ocular tobramycin are localist ocular toxicity and hypersensitivity, including lid itching and swelling and conjunctival erythema. If topical ocular tobramycin is administered concomitantly with systemic amino glycoside antibiotics care should be taken to monitor the total serum concentration. Prolonged use of tobramycin may lead to skin sensitization or result in overgrowth of non susceptible organisms, including fungi. If super infection occurs, appropriate therapy should be initiated. The reactions due to the steroid component in decreasing order of frequency are elevation of intra ocular pressure (IOP) with possible development of glaucoma and infrequent optic nerve damage, posterior subcapsular cataract formation and delayed wound healing.

Manufactured by:

ISIS Pharmaceutical & Chemical Works

25/1-3, Sector 12-C, North Karachi,
Industrial Area, Karachi.

For CIBA Pharmaceutical (Pvt) Ltd.

A-371, Nooriabad, S.I.T.E.,

Karachi-Hyderabad, Super Highway.

سی۔ ٹوب۔ ڈی

۵ ملی لیٹر (تقریباً)

آنکھوں کے قطرے

ٹوبرامائین ۳ ملی گرام (یو ایس پی)

ڈیکسامیٹھاسون ۱ ملی گرام (یو ایس پی)

PRECAUTIONS:

A Steroid glaucoma may be produced after a week or more of treatment in patients predisposed to chronic simple glaucoma. Topical corticosteroid therapy frequently induces intraocular hypertension in normal eyes and increases pressure in eyes with initially elevated pressure. Glaucoma has not always been reversible on cessation of corticosteroid treatment. The local administration of corticosteroid to the eyes of patients with bacterial, viral and fungal conjunctivitis may mask evidence of progression of infection until sight is lost. Corticosteroids may cause progression of the dendritic keratitis (herpes simplex infection), resuming in irreversible clouding of the cornea.

Secondary Infection: Prolonged use of steroids may suppress the host response and thus increase the hazard of Secondary ocular infection. Fungal infections of the cornea are particularly prone to develop coincidentally with long term application of steroid. The possibility of fungal invasion must be considered in any persistent corneal ulceration where steroid treatment has been used. Secondary bacterial ocular infection following suppression of host responses also occurs.

Pregnancy:

The safety of tobramycin during pregnancy has not yet been established.

Nursing Mothers:

It is not known whether this drug is excreted in human milk, therefore caution should be observed when it is administered to a nursing mother.

KNOWN SYMPTOMS OF OVER DOSAGE AND PARTICULARS OF ITS TREATMENT:

See the side effects and special precautions above. Discontinue use immediately.

Presentation:

It is available in sterile dropper bottle of 5 ml Sterile Ophthalmic suspension.

Pharmaceutical precautions:

Store in a cool and dry place, away from light.

Keep out of reach of children.

Storage:

Store below 25°C.

Close bottle after use.

Discard remaining contents four weeks after opening.

خوراک:

اے ۲ قطرے دن میں ۳ سے ۴ مرتبہ ہر ۴ گھنٹہ آنکھوں میں ڈالیں یا ڈاکٹر کی ہدایات کے مطابق استعمال کریں۔

ہدایات:

دوا کو خوب ہلکی، روشنی سے بچائیں۔

۲۵ ڈگری سینٹی گریڈ سے کم درجہ حرارت پر رکھیں۔

تمام ادویات بچوں کی پہنچ سے دور رکھیں۔

انتباہ:

بڑے بچوں کے لئے بعددوا کو ایک ماہ تک استعمال کیا

جاسکتا ہے۔

صرف آنکھوں کے استعمال کیلئے



Marketed by:

LUMEN
PHARMA
www.lumenpharma.com