

CIBAPAT

5ml Approx.
Eye Drops

(Olopatadine 2mg)
Sterile Ophthalmic solution
(USP Specification)

Composition:

Each ml contains:
Olopatadine hydrochloride equivalent to Olopatadine 2.0mg
(USP Specification)

CATEGORY:

Antihistaminic, mast cell stabilizer.

Pharmacodynamics:

Olopatadine is a potent selective antihistaminic and mast cell stabilizing agent that exerts its effects through multiple distinct mechanism of actions. It antagonizes histamine (the primary mediator of allergic response in humans) and prevents histamine induced inflammatory cytokine production by human conjunctival epithelial cells. Olopatadine is a selective H1 receptor antagonist that inhibits Type 1 hypersensitivity reactions. It has been shown to inhibit the release of pro-inflammatory mediators from human conjunctival mast and epithelial cells.

Pharmacokinetics:

The half life in plasma is approximately 8-12 hours and elimination is predominantly through renal excretion approximately 60-70% of the dose is recovered in the urine as parent drug.

Indications:

CIBAPAT (Olopatadine) ophthalmic solution is indicated for the treatment of the signs and symptoms of allergic conjunctivitis.

Dosage and administration:

- The recommended dose of CIBAPAT (Olopatadine) is 1 to 2 drop in each affected eye once a day.
- Place the dropper directly over eye and instill the prescribed number of drops.
- Look downward and gently close your eye for 1 to 2 minutes.
- Place one finger at the corner of the eye near the nose and apply gentle pressure. This will prevent the medication from draining away from the eye.
- Try not to blink and do not rub the eye.
- Do not wear contact lenses while you are using the medicine.

Contraindications:

Olopatadine hydrochloride ophthalmic solution is contraindicated in persons with a known hypersensitivity to olopatadine hydrochloride or any component of the formulation.

Warning & precautions:

Only for Ophthalmic use. Not for injection or oral use. Instillation of CIBAPAT (Olopatadine) may cause transient blurring of vision. Caution should be taken when driving or operating machinery if blurred vision is experienced.

Pregnancy:

There are no adequate and well-controlled studies in pregnant women. CIBAPAT (Olopatadine) should be used in pregnant women only if the potential benefit to the mother justifies the potential risk of the embryo or fetus.

Manufactured by:

ISIS Pharmaceutical & Chemical Works

25/1-3, Sector 12-C, North Karachi,

Industrial Area, Karachi.

For CIBA Pharmaceutical (Pvt) Ltd.

A-371, Nooriabad, S.I.T.E.,

Karachi-Hyderabad, Super Highway

سیلابیٹ
5ml تقریباً
آنکھوں کے قطرے

اولوپاٹاڈین ۲ ملی گرام

Pediatric Use:

Safety and effectiveness in pediatric patient below the age of 3 years have not been established.

Nursing Mothers:

Olopatadine has been identified in the milk of nursing rats following oral administration. It is not known whether topical ocular administration could result in sufficient systemic absorption to produce detectable quantities in the human breast milk. Nevertheless, caution should be exercised when olopatadine is administered to nursing mothers.

Adverse Effects:

Headache has been reported at an incident of 7%. The following undesirable effects have been reported in less than 5% of patients. Asthenia, Blurred vision, Burning or Stinging, Cold syndrome, Dry eye, Foreign body sensation, Hyperemia, Hypersensitivity, Keratitis, Lid edema, Nausea, Pharyngitis, Pruritus, Rhinitis, Sinusitis, and Taste perversion. Some of these events were similar to the underlying disease being studied.

Storage and precaution:

- Store in a cool & dry place, below 25°C.
- Protect from heat and sunlight.
- Keep out of reach of children.
- Discard container 30 days after opening.
- Do not freeze.
- Uses only on medical advice.

Presentation:

CIBAPAT (Olopatadine) Eye drops are available as 5ml sterile dropper bottle.

خوراک:

اسے ۲ قطرے دن میں ایک مرتبہ تھوڑا کچھ آنکھوں میں ڈالیں یا ڈاکٹر کی ہدایات کے مطابق استعمال کریں۔

ہدایات:

دوا کو دھوپ، نمی، روشنی سے بچائیں۔
۲۴ گھنٹہ پہلے سے کم درجہ حرارت پر رکھیں۔
تمام ادویات بچوں کی پہنچ سے دور رکھیں۔

انتباہ:

بچوں کو نلکے کے بعد دوا کو ایک ماہ تک استعمال کیا جاسکتا ہے۔

صرف آنکھوں کے استعمال کیلئے

Marketed by:
 LUMEN
PHARMA
Vision for Life